

Evaluation of Nitrate Reductase Assay For Rapid Detection of Drug Resistant Tuberculosis

¹Ranjit Kumar Sah, ¹Dwij Raj Bhatta, ²Gokarna Raj Ghimire, ³Bhuwaneswor Prasad Kandel,
⁴Bishnu Raj Tiwari, ⁵Jeevan Bahadur Sherchand

¹Central Department of Microbiology, Tribhuvan University, Kirtipur, Kathmandu, Nepal

²National Tuberculosis Center, Thimi, Bhaktapur, Nepal

³Janapriya Multiple Campus, Pokhara, Nepal

⁴School of Health and Allied Sciences, Pokhara University, Pokhara, Nepal

⁵Institute of Medicine, Tribhuvan University Teaching Hospital, Kathmandu, Nepal

ABSTRACT

Emergence of multidrug-resistant tuberculosis (MDR-TB) urgently demands for simple, rapid and inexpensive methods of its detection for the effective treatment of drug resistant tuberculosis, particularly in low-income countries. This prospective study was carried out at National Tuberculosis Reference Laboratory and South Asian Association of Regional Cooperation (SAARC) Tuberculosis and HIV/AIDS Centre, Thimi, Bhaktapur, Nepal, from November 2009 to May 2010 to evaluate nitrate reductase assay (NRA) efficacy for rapid determination of streptomycin, isoniazid, rifampicin and ethambutol susceptibility of Mycobacterium tuberculosis strains. A total of 113 clinical isolates of M. tuberculosis were tested for four first line antitubercular drugs by nitrate reductase assay and were compared with standard proportion method. Results were available in 7-14 days by NRA as compared to proportion method which generally takes 4-6 weeks. The sensitivity and specificity of NRA were 98.1% and 100% for isoniazid, 95.1% and 98.6% for rifampicin, 91.4% and 94.9% for streptomycin, and 78.6% and 97.9% for ethambutol respectively. Agreement between NRA and proportion method were 99.1%, 97.3%, 93.8%, 95.6% for isoniazid, rifampicin, streptomycin and ethambutol, respectively. NRA is easier, inexpensive and reliable method for susceptibility testing of Mycobacterium tuberculosis for isoniazid and rifampicin, the two most important drugs for the treatment of tuberculosis. The reduction in susceptibility testing time, and higher sensitivity and specificity of NRA method is of fundamental importance in detecting MDR-TB.

Key words: Drug susceptibility, MDR-TB, NRA, proportion method

Corresponding address: Ranjit Kumar Sah, Central Department of Microbiology, Tribhuvan University, Kirtipur, Nepal.
E-mail: ranjitkmrsah@gmail.com

INTRODUCTION

Emergence of multidrug resistant tuberculosis (MDR-TB) during the past ten years in higher rate represents a major public health problem, especially in low-resource countries where the burden of the disease is higher. MDR-TB constitutes a serious threat for the effective control of the disease stressing the need for the rapid detection of drug resistance.¹ Conventional methods like proportion method (PM), the resistance ratio method and the absolute concentration method to detect drug resistance in *Mycobacterium tuberculosis* have traditionally relied on slow and cumbersome procedures requiring a minimum of 3–4 weeks to produce results.² Other methods, such as the BACTEC 460 TB System,³ and oxidation-reduction dyes, e.g. tetrazolium⁴, and Microplate Alamar Blue assay (MABA)⁵, are faster but have the drawback of requiring either radioactive or expensive substrates, and are consequently not feasible in most resource-poor settings. A cost effective and rapid drug susceptibility method is required to guide TB treatment.

With similar characteristics to colorimetric methods, the nitrate reductase assay (NRA) has been described, which is

based on the ability of *M. tuberculosis* to reduce nitrate to nitrite after growth in the presence or absence of antibiotics. The use of specific reagents produces a change of color in the presence of nitrites indicating a positive result.⁶ Since it is simpler, cheaper and rapid, evaluation of NRA as an alternative method for determining *M. tuberculosis* susceptibility to rifampicin (RIF), isoniazid (INH), streptomycin (STR) and ethambutol (EMB) is necessary.

MATERIAL AND METHODS

The NRA was performed on a total of 113 isolates of *M. tuberculosis*. Eighty three isolates were from cases reported for pulmonary TB at National Tuberculosis Center (NTC), Thimi, Bhaktapur, Nepal, between November 2009 and May 2010; and thirty isolates were of proficiency testing strain from Supranational Reference Laboratory, Germany. The PM was carried out on Lowenstein and Jenson (LJ) medium according to the standard procedures with the recommended critical concentrations of 40µg/ml for rifampicin, 0.2µg/ml for isoniazid, 2µg/ml for ethambutol and 4µg/ml for streptomycin.² NRA was performed according to the already

described protocol.⁶ The following critical concentrations were used: 0.2µg/ml for INH, 40 µg/ml for RIF, 4µg/ml for STR and 2.0µg/ml for EMB. Briefly, fresh subculture (1µl loops of bacteria) from isolates of *M. tuberculosis* grown on LJ medium was taken and vortexed in small amount of sterile distilled water and turbidity was adjusted according to McFarland standard no.1. Part of the suspension was diluted 1:10 in sterile distilled water. For each isolate, 0.2 ml of suspension was inoculated into the tubes containing LJ medium with potassium nitrate (KNO₃) and the antitubercular drugs; 0.2 ml of the 1:10 dilution was inoculated into drug free media (LJ media) containing KNO₃ which served as growth controls. Tubes in triplicate were incubated at 37°C for 14 days and 0.5 ml of a mixture of three reagents (25µl of concentrated HCl, 50µl of 2% sulphanilamide and 50µl of 1% n-1-naphthyl-ethylenediamine dihydrochloride) was added to one drug-free control tube after 7 days of incubation. If its colour changed to pink then tubes with drugs were tested. An isolate was considered resistant if there was colour change (pink or deep red to violet) in the drug tube in question greater than in the 1:10 diluted growth control on the same day. If the tubes did not show any colour change and remains the same, these were further incubated for 10 days and for 14 days as described.⁶ Statistical analysis of data was carried out using χ^2 test at 5% level of significance using Statistical Package for Social Science (SPSS version 17.0).

RESULTS

Altogether 113 *M. tuberculosis* isolates were tested and the results were available in 7 days for 63.3% of the strains, in 10 days for 82.3% of the strains, and in 14 days for 100% of the strains by NRA method (Table 1). Of the total isolates, isolates showing resistance to INH, RIF, STR and EMB were 46.0%, 36.3%, 30.9% and 12.4% respectively, by the proportion method while resistance to respective drugs was 45.1%, 35.4%, 31.8% and 11.5% by NRA. The sensitivity and specificity for NRA were 98.1% and 100.0%, 95.1% and 98.6%, 91.4% and 94.9%, and 78.6% and 97.9% for INH, RIF, STR and EMB respectively (Table 2). The results showed that NRA and proportion method do not differ significantly ($P>0.05$ for all the drugs). There was high agreement between both methods (Table 3) when tested against INH, RMP, SM and EMB with kappa, $k=0.98, 0.93, 0.86$ and 0.81 respectively.

Table 1: Number of days required for result by NRA method

No. of days	No. of specimens reported	Cumulative %
7	56	63.3
10	37	82.3
14	20	100.0

Table 2: Comparison of indirect nitrate reductase assay results with conventional proportion method

Drugs	Conventional Proportion method	Nitrate reductase assay method					
		Resistant	Sensitive	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)
INH	Resistant=52	51	1	98.1	100	100	98.4
	Sensitive=61	0	61				
RFP	Resistant=41	39	2	95.1	98.6	97.5	97.3
	Sensitive=72	1	71				
SM	Resistant=35	32	3	91.4	94.9	88.9	96.1
	Sensitive=78	4	74				
EMB	Resistant=14	11	3	78.6	97.9	84.6	97
	Sensitive=99	2	97				

Table 3: Percentage agreement between the proportion and the NRA methods for susceptibility testing of *M. tuberculosis* to each drug tested

S.N.	Drugs	No. of isolates with the following results		Percent agreement
		PR method-Susceptible	PR method-Resistant	
		NRA method-usceptible	NRA method-Resistant	
1	INH	61	51	99.1
2	RFP	71	39	97.3
3	SM	74	32	93.8
4	EMB	97	11	95.6

DISCUSSION

Traditional drug susceptibility testing such as the PM on LJ or agar medium is time consuming. For developing countries, it would be helpful to have a simple and inexpensive test that can rapidly detect resistant *M. tuberculosis* strains. The reporting time of NRA was between 7-14 days with majority in 10 days as against 28-42 days for the conventional PM. This is very much comparable with the results of MGIT and BACTEC 460TB, which require the use of expensive instruments and high running cost. The rapidity with which test could be performed has also been reported with comparable accuracy to that of NRA such as the MTT or resazurin assays.⁷ However, they make use of liquid medium in a micro-plate format and that makes the technique more complex and might also constitute a biohazard. Instead, the NRA utilizes standard solid LJ-medium, although with KNO₃ incorporated and it could therefore be easily adopted in any culture laboratory.

Nitrate reductase-negative strains of *M. tuberculosis* are very unusual⁸ and on the other hand, false susceptible results would in this case be detected by the lack of a positive reaction also in drug free growth. *M. bovis* does not reduce nitrate, therefore the NRA technique is not applicable. Disadvantages of NRA are; the culture is killed by the mix reagent used to develop the assay, requiring that multiple cultures be prepared if comparative testing will be performed and only fresh cultures must be used (<14 days).

In present study, a high level of agreement between NRA and PM was determined for isoniazid and rifampicin. However, despite a higher agreement between NRA and PM for streptomycin (93.8%) and ethambutol (95.6%), low sensitivity of streptomycin (91.4%) and ethambutol (78.6%) was found. For isoniazid and rifampicin, sensitivity and specificity was found to be high; 98.1% and 100.0%, and 95.1% and 98.6% respectively. These results are very important since rifampicin and isoniazid are the two most important drugs used in the treatment of TB.⁹ Higher accuracy results for INH, RIF and EMB and lower accuracy results for STR were shown in a similar multicenter study.¹⁰ The percentage agreement of 98.3%, 98.3%, 90.8% and 93.3% by NRA for INH, RMP,

SM and EMB respectively, were observed in Brazilian study.¹¹ However, marginally lower sensitivity and specificity for STR and EMB have also been reported.^{7,12} For STR and EMB testing, further studies are required to optimize preparation of adequate inoculum and drug concentration in order to avoid strain misclassification.

CONCLUSION

In the context of emergence of MDR-TB, the NRA may be of great importance due to its higher sensitivity and specificity for the rapid detection of rifampicin and isoniazid resistance, the two most important drugs for tuberculosis treatment.

ACKNOWLEDGEMENT

The authors acknowledge all the laboratory staffs of NTC and SAARC TB/HIV Center, for their worthy cooperation during the study.

REFERENCES

1. Caminero JA. Multidrug-resistant tuberculosis: epidemiology, risk factors and case finding. *Int J Tuberc Lung Dis.* 2010; 14(4):382-390.
2. Canetti G, Fox W, Khomeiko A, Mahler HT, Menon NK, Mitchison DA, Rist N, Smelev NA. Advances in techniques of testing mycobacterial drug sensitivity and the use of sensitivity tests in tuberculosis control programmes. *Bull World Health Org.* 1969; 41:21-43.
3. Roberts GD, Goodman NL, Heifets L, Larsh HW, Lindner TH, McClatchy JK, McGinnis MR, Siddiqi SH, Wright P. Evaluation of the BACTEC radiometric method for recovery of mycobacteria and drug susceptibility testing of *Mycobacterium tuberculosis* from acid-fast smear-positive specimens. *J Clin Microbiol.* 1983;18:689-96.
4. Caviedes L, Delgado J, Gilman RH. Tetrazolium microplate assay as a rapid and inexpensive colorimetric method for determination of antibiotic susceptibility of *Mycobacterium tuberculosis*. *J Clin Microbiol.* 2002; 40:1873-74.
5. Franzblau SG, Witzig RS, McLaughlin JC, Torres P, Madico G, Hernandez A, Degnan MT, Cook MB, Quenzer VK, Ferguson RM, Gilman RH. Rapid, low-technology MIC determination with clinical *Mycobacterium tuberculosis* isolates by using the microplate Alamar Blue assay. *J Clin Microbiol.* 1998; 36:362-66.
6. Angeby KA, Klintz L, Hoffner SE. Rapid and inexpensive drug susceptibility testing of *Mycobacterium tuberculosis* with a nitrate reductase assay. *J Clin Microbiol.* 2002; 40:553-55.
7. Montoro E, Lemus D, Echemendia M, Martin A, Portaels F, Palomino JC. Comparative evaluation of the nitrate reduction assay, the MTT test, and the resazurin microtitre assay for drug susceptibility testing of clinical isolates of *Mycobacterium tuberculosis*. *J Antimicrob Chemother.* 2005; 55:500-505.
8. Kent PT, Kubica GP. *Public health mycobacteriology: A guide for the level III laboratory.* U.S. Department of Health and Human Services. Centers for Disease Control and Prevention. Atlanta, Georgia, USA. 1985; 159-84.
9. World Health Organization (WHO). The WHO/International Union against Tuberculosis and Lung Disease. Global Project on Anti-tuberculosis Drug Resistance Surveillance. *Antituberculosis Drug Resistance in the World*, report no. 3. World Health Organization. Geneva, Switzerland. 2004.
10. Martin A, Montoro E, Lemus D, Simboli N, Morcillo N, Velasco M et. al. Multicenter evaluation of the nitrate reductase assay for drug resistance detection of *Mycobacterium tuberculosis*. *J Microbiol Methods.* 2005; 63:145-50.
11. Shikama Mde L, Silva RR, Martins MC, Giampaglia CM, Oliveira RS, Silva RF et. al. Rapid detection of resistant tuberculosis by nitrate reductase assay performed in three settings in Brazil. *J Antimicrob Chemother.* 2009; 64:794-96.
12. Lemus D, Montoro E, Echemendia M, Martin A, Portaels F, Palomino JC. Nitrate reductase assay for detection of drug resistance in *Mycobacterium tuberculosis*: simple and inexpensive method for low-resource laboratories. *J Med Microbiol.* 2006; 55:861-63.